

On the efficiency of nonlocal gates generation

Berry Groisman^a and Benni Reznik^b

^a*HH Wills Physics Laboratory, University of Bristol, Tyndall Avenue, Bristol, BS8 1TL, UK*

^b*School of Physics and Astronomy, Raymond and Beverly Sackler Faculty of Exact Sciences, Tel Aviv University, Tel Aviv 69978, Israel*

We propose and study a method for using non-maximally entangled states to implement probabilistically non-local gates. Unlike distillation-based protocols, this method does not generate a maximally entangled state at intermediate stages of the process. As a consequence, the method becomes more efficient at a certain range of parameters. Gates of the form $\exp[i\xi\sigma_{n_A}\sigma_{n_B}]$ with $\xi \ll 1$, can be implemented with nearly unit probability and with vanishingly small entanglement, while for the distillation-based method the gate is produced with a vanishing success probability. We also derive an upper bound to the optimal success probability and show that in the small entanglement limit, the bound is tight.

I. INTRODUCTION

Over the recent years it has been shown that entanglement can be used to perform various quantum processes such as teleportation, quantum communication, quantum cryptography and quantum computation [1]. One important use of entanglement, involves the implementation of quantum gates (or more generally, interaction) between spatially separated qubits, without actually having to transport the physical system that carries the state from one place to another. The study of such “non-local” gate operations, has some bearing on practical and fundamental issues. On one hand, the connection between entanglement as a resource for non-local gates, and the related problem of quantifying the capability of non-local gates to generate entanglement, deals with the fundamental relation between entanglement and interactions. This problem has been studied by several groups [2, 3, 4, 5, 6, 7, 8, 9] but is yet not fully understood. On the other hand, non-local gates may be used as primitives in protocols involving several separated systems, for instance, in multi-party computation problems.

In principle, any non-local gate can be implemented given by sufficient amount of shared entanglement, and by exchanging a sufficient number of classical bits: we can use quantum teleportation to teleport the relevant qubits states to a single location, apply *locally* the relevant interaction in order to generate the desired gate, and finally teleport the states back to their original position. In general however, the implementation of gates with quantum teleportation methods may not be efficient, and provides only an upper bound on the required amount of entanglement. Since entanglement is an expensive resource, it is important to optimize its usage and search for economic methods for implementing nonlocal gates.

In this paper we shall reconsider the problem of gates acting on two qubits which have the structure

$$U_{AB}(\xi) = e^{i\xi\sigma_{n_A}\sigma_{n_B}}. \quad (1)$$

This family of gates includes the controlled-NOT (CNOT) gate, which up to local rotations corresponds to $U_{AB}(\xi = \pi/4)$. By using teleportation, $U_{AB}(\xi)$ can




FIG. 1: A non-local gate can be implemented using distillation-based method which starts by generating a maximally entangled state in an intermediate stage of the protocol, or by a more direct method that avoids distillation.

be implemented for any ξ using two maximally entangled pairs (e-bits). It was shown however that, if n_A and n_B are locally known, one e-bit is sufficient [5, 6]. Moreover, it was recently shown that for the cases $\xi = \pi/2^N$, where $N > 2$, this gate can be implemented with less than one e-bit [7]. The later method utilizes N pairs, each carrying a different amount of entanglement, that sums up to less than one e-bit.

In this article we shall study the possibility of realizing the gate (1) using a *single* copy of a *non-maximally* entangled state

$$|\phi_{ab}(\alpha)\rangle = \cos \alpha |0_a\rangle |0_b\rangle + \sin \alpha |1_a\rangle |1_b\rangle, \quad (2)$$

In this case, as we show in the sequel, with certain states the gate can not be implemented deterministically [10]. Hence in the rest of the paper we shall study methods for probabilistic implementations.

The simplest method to generate (1) is offered by distillation methods. To this end, one can first transform with some probability the state (2) to a the maximally entangled pair

$$|\phi_{ab}^+\rangle = \frac{1}{\sqrt{2}}(|0_a\rangle |0_b\rangle + |1_a\rangle |1_b\rangle), \quad (3)$$

and then utilizes a deterministic scheme (e.g. that suggested in [6] in order to implement the gate (see Fig. (1)). In this scheme, the probability to implement the gate successfully is determined by the probability, p_{dis} to convert the non-maximally entangled pair (2) into maximally entangled pair (3). It was recently shown that

the maximal value of p_{dis} is twice the modulus square of the Schmidt coefficient of smaller magnitude [11, 12], i.e. $p_{dis} = 2(\sin \alpha)^2$ and $0 < \alpha < \pi/4$. The simple method to distill an e-bit out of a non-maximally entangled state is known as the Procrustean method [13].

We notice that the success probability of a distillation-based method does not depend on the gate parameter ξ , and decreases with α . This suggests that in a more efficient method one could optimize the scheme as a function of the gate parameter ξ as well. As we show, an improvement of the distillation-based protocols can be indeed obtained when the process does not require in an intermediate step a maximally entangled state (Fig. 1).

To see that this is possible, consider the special case of $\alpha = \xi$. Let Alice and Bob start by performing local CNOT interactions: Alice applies CNOT (with respect to σ_{n_A}) between her qubits a and A , described by the unitary transformation

$$U_{aA} = |0_a\rangle\langle 0_a| \otimes I_A + |1_a\rangle\langle 1_a| \otimes \sigma_{n_A}. \quad (4)$$

Bob does the same on his side. This yields the four-particle state

$$\left(\cos \alpha |0_a 0_b\rangle \otimes I_{AB} + \sin \alpha |1_a 1_b\rangle \otimes \sigma_{n_A} \sigma_{n_B} \right) |\Psi_{AB}\rangle \quad (5)$$

Next they measure a and b : Alice measures σ_y and Bob measures σ_x . The result can be then written as

$$\left(\cos \alpha I_{AB} \pm i \sin \alpha \sigma_{n_A} \sigma_{n_B} \right) |\Psi_{AB}\rangle = e^{\pm i \alpha \sigma_{n_A} \sigma_{n_B}} |\Psi_{AB}\rangle, \quad (6)$$

where the $+$ ($-$) sign is obtained with probability $1/2$ when the measurement results satisfies $\sigma_x = \sigma_y$ ($\sigma_x = -\sigma_y$). Hence we showed that the success probability for gates with $\alpha = \xi$ is here given by $1/2$! which can be larger compared with p_{dis} for small enough values of α .

At first sight this result seems paradoxical [14]. Recall the entanglement capability the gate (1) is known in this limit to be $\sim \xi$ [8]. Hence we have started with an entanglement proportional to $\xi^2 = \alpha^2$ and ended up with more entanglement $\sim \xi$ with probability $1/2$! The resolution to this paradox, is connected with the fact that the maximally capability is obtained by acting the gate on a partially entangled state. Hence in $1/2$ of the times we end up by *reducing* the entanglement. The net increase is not larger than ξ^2 .

In the rest of this paper we present and study a method for generating gates, that does not involve maximal entanglement states at intermediate steps, and hence utilizes the above mentioned idea of a “direct” map between entanglement and gates (Fig. 1). The method can be used for arbitrary values of ξ and α , and for a certain range of parameters, is more efficient compared with the distillation-based (e.g. Procrustean) methods. In particular, we show that gates with very a small angle ξ , can be implemented with probability very close to unity, with vanishingly small entanglement. This is to be compared with the distillation-based method, which in this case succeeds with a vanishingly small probability.

We shall also exam the general relation between the maximal success probability p_{max} , the gate parameter ξ , and the given entangled state, described by the parameter α .

The article proceeds as follows. In the next section II we derive a bound on p_{max} for a certain range of parameters ξ and α . In Sec. III we present a “direct” method for probabilistic generation of non-local gates. Then, in Sec. IV we refer to one of these gates as a target gate and analyze the probability of its implementation. Based on these results we show that the bound derived in Sec. II is tight.

II. A BOUND ON THE SUCCESS PROBABILITY

In this section we derive an upper bound on the maximal probability of successful implementation of the gate (1) on the state $|\Psi_{AB}\rangle$ using (2) as a resource when $\alpha < \xi$.

We can write a required task as a transformation

$$(\cos \alpha |0_a\rangle\langle 0_b| + \sin \alpha |1_a\rangle\langle 1_b|) |\Psi_{AB}\rangle \xrightarrow{p} e^{i \xi \sigma_{n_A} \sigma_{n_B}} |\Psi_{AB}\rangle = (\cos \xi + i \sin \xi \sigma_{n_A} \sigma_{n_B}) |\Psi_{AB}\rangle, \quad (7)$$

where the p denotes the probability of the above transition.

Let us consider the special case of the product state

$$|\Psi_{AB}\rangle = |\uparrow_{n_A}^\perp\rangle |\uparrow_{n_B}^\perp\rangle,$$

where the axis $n^\perp$ are such that $\sigma_n |\uparrow_{n^\perp}\rangle = |\downarrow_{n^\perp}\rangle$. In this case

$$(\cos \alpha |0_a\rangle\langle 0_b| + \sin \alpha |1_a\rangle\langle 1_b|) |\uparrow_{n_A}\rangle |\uparrow_{n_B}\rangle \xrightarrow{p} \cos \xi |\uparrow_{n_A}\rangle |\uparrow_{n_B}\rangle + i \sin \xi |\downarrow_{n_A}\rangle |\downarrow_{n_B}\rangle \quad (8)$$

The main idea is to use the *majorization condition* [11, 12, 17] in order to bound the probability p . If $\alpha < \xi$, then $\{(\cos \alpha)^2, (\sin \alpha)^2\}$ does not majorized by $\{(\cos \xi)^2, (\sin \xi)^2\}$, which means that the transformation (8) cannot be performed with certainty. The maximal probability of this transformation given in this case by

$$p_{max} = \frac{(\sin \alpha)^2}{(\sin \xi)^2}. \quad (9)$$

Now we apply to the general result from [15] that shows that the probability of successful simulation of a unitary is independent of the input state. Thus, the bound (9) obtained for the product state is as good as a bound one might obtain for all other $|\Psi_{AB}\rangle$. In our particular case it can be also easily checked explicitly that any other $|\Psi_{AB}\rangle$ will give the same upper bound on p .

In the Sec. IV we will show that for $\alpha \ll \xi$ this bound is tight.

In passing we note that for $\xi = \pi/4$ the bound (9) is consistent with the results of [15, 16] concerning an optimal simulation of non-local CNOT gate, i.e. $U_{AB}(\xi = \pi/4)$, using $U_{AB}(\xi < \pi/4)$.

We note that for $\alpha \geq \xi$ the majorization method does not provide any restrictions on the transformation (8), i.e. we cannot use it to derive the bound in this case. The majorization does not prevent p_{max} even from reaching one when $\alpha = \xi$, although for small ξ this possibility can be discarded based on the results on the entanglement capability [8]. It is natural to conjecture that in the single state case, $p_{max} = 1$ may be achieved only for $\alpha = \pi/4$. It is unlikely that p_{max} exhibits non-smooth behavior. Nevertheless, the question whether the gate (1) can be generated deterministically for $\alpha \geq \xi$ is still open.

III. A METHOD FOR PROBABILISTIC GENERATION OF NON-LOCAL GATES

In this section we demonstrate how Alice and Bob can probabilistically generate a pair of gates of the type (1) directly from $|\phi_{ab}(\alpha)\rangle$ without distilling it.

A. Mapping states to Stator

In order to simplify the explanation of our method we use a hybrid state-operator object (stator) defined in [6]. The Stator describes quantum correlations between the state of one systems and the operation/s acting on another system/s.

How do we prepare the stator? Alice and Bob start by performing the first step as described in Sec. I - Eq. (4) and (5).

Next Bob performs a measurement of σ_x of the entangled qubit b to project out a certain value. The resulting state is now

$$(|0_b\rangle \pm |1_b\rangle) \otimes \left(\cos \alpha |0_a\rangle \otimes I_B \pm \sin \alpha |1_a\rangle \otimes \sigma_{n_A} \sigma_{n_B} \right) |\Psi_{AB}\rangle \quad (10)$$

Finally Bob informs Alice what was the result of his measurement by sending Alice one classical bit of information. For the case that $\sigma_x = -1$ Alice performs a trivial π rotation around the $\hat{z}$ axis and flips the $-$ sign to a $+$ sign. The resulting state of the system is now given by

$$(|0_b\rangle \pm |1_b\rangle) \otimes \left(\cos \alpha |0_a\rangle \otimes I_B + \sin \alpha |1_a\rangle \otimes \sigma_{n_A} \sigma_{n_B} \right) |\Psi_{AB}\rangle \quad (11)$$

Since Bob's previously entangled qubit factors out, the final state of Alice's qubits A , a and Bob's qubit B is

$$\left(\cos \alpha |0_a\rangle \otimes I_B + \sin \alpha |1_a\rangle \otimes \sigma_{n_A} \sigma_{n_B} \right) |\Psi_{AB}\rangle \equiv S |\Psi_{AB}\rangle, \quad (12)$$

where the stator S captures the correlation between the state $|0_a\rangle$ and $|1_a\rangle$ of Alice's qubit (a) and unitary transformation I_{AB} and $\sigma_{n_A} \sigma_{n_B}$ acting on $|\Psi_{AB}\rangle$.

If $\alpha = \pi/4$, then the stator is "maximal" and can be used to apply (1) with certainty [6]. This is done by utilizing the identity:

$$e^{i\xi\sigma_{x_a}} S = S e^{i\xi\sigma_{n_A} \sigma_{n_B}}. \quad (13)$$

Hence by applying a local rotation by Alice we can "pull out" the required unitary operator that generates the gate. In the following section we show how to use "non-maximal" stator in order to apply (1) probabilistically.

B. Generating non-local gates

Our goal is to find an appropriate $\theta(\xi, \alpha)$ such that by applying locally a unitary on S one gets the sum of probabilistic unitaries

$$e^{i\theta(\xi, \alpha)\sigma_{x_a}} S = \sqrt{p} |0_a\rangle \otimes e^{i\xi\sigma_{n_A} \sigma_{n_B}} + \sqrt{1-p} |1_a\rangle \otimes \sigma_{n_A} \sigma_{n_B} e^{i\tilde{\xi}\sigma_{n_A} \sigma_{n_B}}. \quad (14)$$

As a result the state of Alice's particle a will be correlated with two different gates applied on the state $|\Psi\rangle_{AB}$: if Alice will measure her particle a in z -basis, then she will get $|0_a\rangle$ with probability p and the nonlocal gate $U_{AB}^{(1)} = \exp(i\xi\sigma_{n_A} \sigma_{n_B})$ will be generated. Similarly, when she gets $|1_a\rangle$ (that happens with probability $1-p$) then the nonlocal gate $U_{AB}^{(2)} = \exp(i\tilde{\xi}\sigma_{n_A} \sigma_{n_B})$ is generated (up to trivial local rotations $\sigma_{n_B} \sigma_{n_B}$).

It is straightforward to verify that Alice has to apply in (14) the unitary with

$$\theta(\xi, \alpha) = \tan^{-1} \left[\frac{\tan \xi}{\tan \alpha} \right] \quad (15)$$

and to show that ξ , $\tilde{\xi}$, p , and α have to satisfy the following relations:

$$p = (\cos \alpha \cos \theta)^2 + (\sin \alpha \sin \theta)^2, \quad (16)$$

$$\xi = \cos^{-1} \left[\frac{\cos \alpha \cos \theta}{\sqrt{p}} \right] \quad (17)$$

$$\tilde{\xi} = \cos^{-1} \left[\frac{\sin \alpha \cos \theta}{\sqrt{1-p}} \right] \quad (18)$$

$$\frac{\tan \xi}{\tan \tilde{\xi}} = (\tan \alpha)^2 \quad (19)$$

$$p = \frac{(\sin \alpha)^2}{1 - \frac{1-2(\sin \alpha)^2}{1-(\sin \alpha)^2} \cos^2 \xi} \quad (20)$$

Thus, as it follows from (19), given a nonmaximally entangled pair (2) as a resource Alice and Bob can generate any the pair of gates $U_{AB}^{(1)}, U_{AB}^{(2)}$ (each one with appropriate probability) where two parameters $\xi, \tilde{\xi}$ which satisfy (19). Since for fixed α infinitely many pairs of angles $\xi, \tilde{\xi}$ (in the range $\pm\pi/2$) which satisfy (19) can be found, infinitely many pairs of gates can be generated.

To obtain the required gate they choose a pair $\xi, \tilde{\xi}$ that satisfies (19), and calculate θ according to (15). Finally Alice applies $U_a = \exp(i\theta\sigma_{x_a})$, and measures the operator σ_z of qubit a .

We note that for the special case of "maximal" S , i.e. when $\alpha = \pi/4$, we obtain $p = \frac{1}{2}$ and $\theta = \xi = \tilde{\xi}$. Thus we get a deterministic gate as in [6].

IV. OPTIMAL GENERATION OF A SINGLE GATE

Let us return to our original problem. Suppose that Alice and Bob are interested to implement a gate with a particular value of ξ . Following the scenario presented in Sec. III Alice and Bob will successfully implement the desired gate ($U_{AB}(\xi)$) with probability p and will fail, i.e. implement the "trash" gate ($U_{AB}(\tilde{\xi})$), with probability $1-p$. In the Fig. 2 the probability of success p as a function of $\sin^2 \alpha$ is plotted for various values of ξ . Below we have list the important features of the resulting behavior:

- Above the threshold $\xi_c = \pi/4$ the Procrustean method is better for all values of α . For $\xi < \pi/4$ our method gives higher probability of success in certain range of α . This range increases as ξ decreases and becomes dominant at small ξ . We note, however, that every gate (1) with $\xi > \pi/4$ can be decomposed into the gate with $|\xi| < \pi/4$ followed by *local* rotations. Thus, the case with $\xi > \pi/4$ is not really interesting from nonlocal point of view.




FIG. 2: The probability of success p as a function of $\sin^2 \alpha$ for various values of ξ . The dashed line shows the probability $2\sin^2 \alpha$ given by the Procrustean method. Points A and C correspond to α_{opt} and α_{opt}^{Proc} for $\xi = \pi/64$. Thus, if the resource entanglement is in the interval (A,B), then Alice and Bob can decrease it to optimal A and achieve p_{max} taking an advantage over the Procrustean method. Point B corresponds to the value $\sin^2 \alpha_c = 1/3$, where both methods give same nonzero probability as $\xi \rightarrow 0$. (The values of ξ presented on the plot are taken to be proportional to $1/2^n$ for demonstration only: our method works for all real values of ξ in the appropriate interval.)

- For each fixed value of ξ there is an *optimal* state (2) with $\alpha = \alpha_{opt}$, where

$$\sin^2 \alpha_{opt} = \frac{\sin^2 \xi + 0.5 \sin 2\xi}{\cos 2\xi}, \quad (21)$$

which gives the maximum probability of success p_{max} .

- For a particular value of ξ our method is obviously better for all $\alpha < \alpha_c$, where

$$\sin^2 \alpha_c = \frac{2 \cos^2 \xi - 1}{4 \cos^2 \xi - 1} \quad (22)$$

is the point where both methods cross. This value goes to $1/3$ at the limit $\xi \rightarrow 0$. One might think, that for $\alpha > \alpha_c$ the Procrustean method is always better. However, this is not the case. Indeed, the Procrustean method achieves the value equal to p_{max} , at some $\alpha_{opt}^{Proc} > \alpha_c$, where the value of α_{opt}^{Proc} obeys the condition:

$$p(\alpha_{opt}) = 2 \sin^2 \alpha_{opt}^{Proc}. \quad (23)$$

Thus, if Alice and Bob are given by the state (2) with $\alpha_{opt} < \alpha < \alpha_{opt}^{Proc}$ then they always can convert it to the optimal state and achieve p_{max} taking




FIG. 3: The p vs. $\sin^2 \alpha$ dependence for $\xi = 0.014$ rad. Two linear regimes $p = 1 - \sin^2 \alpha$ and $p = 5102\alpha^2$ are presented. The maximal probability corresponds to $\alpha_{opt}^2 = 0.014$ and equals to $p_{max} = 0.986$.

an advantage over the Procrustean method, which is better only for α in the range $\alpha_{opt}^{Proc} < \alpha < \pi/4$. In the following we will show that α_{opt}^{Proc} becomes very close to $\pi/4$ in the limit of small ξ .

- For all values of ξ at the limit $\alpha \ll \xi$ the expression (20) can be approximated by a linear (with respect to $\sin^2 \alpha$) dependence

$$p = \frac{(\sin \alpha)^2}{(\sin \xi)^2}, \quad (24)$$

which is consistent with the bound (9), i.e. this bound is tight at this limit.

- At the limit $\xi \rightarrow 0$ an additional linear regime is obtained: if $\xi \ll \alpha$ then p is approximated by

$$p = 1 - \sin^2 \alpha. \quad (25)$$

The maximal probability is obtained at $\alpha_{opt}^2 = \xi$:

$$p_{max} = 1 - \alpha_{opt}^2 = 1 - \xi \quad (26)$$

Fig. 3 shows both linear regimes and p_{max} for $\xi = 0.014$ rad. As we can see, at the limit $\xi \rightarrow 0$ the maximal probability of success goes to unity as $p_{max} = 1 - \xi$. Surprisingly, this probability is obtained for very weakly entangled optimal state (2) with $\alpha_{opt}^2 = \xi$.

The Procrustean method achieves this probability of success only for $\sin^2 \alpha \geq (1 - \xi)/2$, i.e. for α which are very close to $\pi/4$. In other words, at the limit of small ξ our method is better than the Procrustean method almost for all resource states (2), except for those that are very close to maximally entangled.

From (19) we can find $\tilde{\xi}$ of the "trash" gate that will be generated in the case of failure for every particular ξ and α . We note that each of two gates possesses certain entanglement capability, i.e. ability to generate certain amount of entanglement. It can be checked explicitly that the corresponding convex sum of the entanglement capabilities of two gates does not exceed the entanglement consumed in the process, i.e. the entanglement possessed by (2).

V. COMPARISON WITH PREVIOUS WORK

In this section we compare our results with previous work on the interconvertibility of two nonlocal operations [15, 16]. These articles use an isomorphism between states and operations, in order to generate a transformation between two non-local gates. In the sense already discussed in the introduction, this method is indirect: $U_1 \rightarrow \psi_1 \rightarrow \psi_2 \rightarrow U_2$. First the given non-local gate U_1 is applied on a standard non-entangled state and gives rise to the entangled state ψ_1 . Then at an intermediate step, ψ_1 is mapped (distilled) to the desired entangled state ψ_2 , and finally the isomorphism is used again to regenerate the desired non-local gate U_2 .

The main motivation of [15, 16] was searching for equivalent classes, i.e. the *possibility* of such a transition, so a probability optimization was not an issue in general - any nonzero probability was good in principle. Thus, the problem of optimizing the probability for interconversion between two non-local gates $U_{AB}(\alpha)$ and $U_{AB}(\xi)$ was not addressed in general, however an answer was given for the special case of $\xi = \pi/4$. In this case, any $U_{AB}(\alpha)$ can be deterministically obtained from $U_{AB}(\pi/4)$, while $U_{AB}(\pi/4)$ can be obtained from $U_{AB}(\alpha)$ with optimal probability $2(\sin \alpha)^2$, that follows from majorization condition used on the second step ($\psi_\alpha \rightarrow \psi_\xi$) and from the fact that $p[\psi_\xi \rightarrow U_{AB}(\xi)] = 1$ in this case. In the more general case, $\alpha < \xi < \pi/4$, however, the optimal probability $p[\psi_\xi \rightarrow U_{AB}(\xi)]$ is not known. Moreover, the results of our present paper suggest that using state-operation isomorphism might be not the most efficient way to address the problem of efficient non-local gate interconversion, due to the loss of efficiency at the intermediate indirect $\psi_\alpha \rightarrow \psi_\xi \rightarrow U_\xi$ transformation. Indeed, for $\alpha < \xi$ our method takes ψ_α directly to $U_{AB}(\xi)$ and achieves the maximal probability (9). If, however, one goes through ψ_ξ at an intermediate stage then the same value of probability is multiplied by $p[\psi_\xi \rightarrow U_{AB}(\xi)]$, which is up to date known to be $1/2$ according to [7] and our present work.

VI. CONCLUSION

In this paper we have presented a method for probabilistic implementation of nonlocal operations (characterized by an interaction parameter ξ) on two qubits us-

ing single nonmaximally entangled state (characterized by an entanglement monotone α) shared by the parties. Our method is characterized by a direct utilization of a nonmaximally entangled pair without having to convert it first to a maximally entangled pair as an intermediate step. We found that if the resource nonmaximally entanglement is below a certain value, our method has a higher success probability than the distillation-based (e.g. Procrustean) methods. This critical value increases with decreasing of ξ , so our method is almost always preferable at small ξ . For very small ξ the probability reaches the value $p_{max} = 1 - \xi$ in contrast to $p = 2\xi$ that is given by the Procrustean method for the same α . The explanation of this efficiency gain seems to be that our method, unlike the Procrustean method, does not have to create maximal entanglement at an intermediate stage of the protocol. On the other hand, one might think that an advantage of the distillation-based methods is that they do not affect the target state in the case of failure. We note, however, that our method changes the state in a *known* way.

Interestingly, we found that for higher success probability our method requires as a resource pairs with

less entanglement. For example, for given α and very small ξ the maximal probability of success is obtained at $\sin^2 \alpha = \xi$. Thus, when a state with a higher then optimal entanglement is given, the parties have to reduce the entanglement to the optimal lower value, before implementing our protocol. This behavior is consistent with the results of [9].

We have also addressed the fundamental question of the theoretical bound on the probability of successful implementation of a nonlocal gate. We found an upper bound for certain range of parameters and showed that this bound is tight. However, the general question of the upper bound for the whole range of parameters remains open.

Acknowledgments

We would like to thank Sandu Popescu for very useful discussions. BG acknowledges the European IST-FET project RESQ. BR acknowledges the ISF, grant 62/01-1.

-
- [1] M.A. Nielsen and I.L.Chuang, Quantum Computation and Quantum Information, Cambridge University Press (2004).
 - [2] J. Eisert, K. Jacobs, P. Papadopoulos, M.B. Plenio, Phys. Rev. A **62**, 052317 (2000); e-print quant-ph/0005101.
 - [3] B. Kraus and J.I. Cirac, Phys. Rev. A **63**, 062309 (2001); e-print quant-ph/0011050.
 - [4] D. Collins, N. Linden and S. Popescu, Phys. Rev. A **64** 032302 (2001); e-print quant-ph/0005102.
 - [5] S. F. Huelga, J. A. Vaccaro, A. Chefles and M. B. Plenio, Phys. Rev. A **63** 042303 (2001); S. Huelga, M.B. Plenio and J.A. Vaccaro, e-print quant-ph/0107110.
 - [6] B. Reznik, Y. Aharonov, and B. Groisman, Phys. Rev. A **65** 032312 (2002).
 - [7] J. I. Cirac, W. Dür, B. Kraus and M. Lewenstein, Phys. Rev. Lett, **86**, 544 (2001); W. Dür and J. I. Cirac, Phys. Rev. A, **64**, 012317 (2001).
 - [8] W. Dür, G. Vidal, J. I. Cirac, N. Linden, and S. Popescu, Phys. Rev. Lett, **87**, 137901 (2001).
 - [9] B. Reznik, e-print quant-ph/0203055.
 - [10] This is always the case when $\alpha < \xi$ (see Sec. II). However, for $\alpha \geq \xi$ the existence of deterministic procedure is an open question.
 - [11] H.-K. Lo, S. Popescu, Phys. Rev. A , **63** 022301 (2001).
 - [12] G. Vidal, Phys. Rev. Lett, **83** 051046 (1999).
 - [13] C.H.Bennett, J.H. Bernstein, S. Popescu, B. Schumacher, Phys. Rev. A , **53** 2046 (1996).
 - [14] As was first pointed out by Sandu Popescu (private communication).
 - [15] W. Dür, G. Vidal, J. I. Cirac, Phys. Rev. Lett. , **89**, 057901 (2002).
 - [16] W. Dür and J. I. Cirac, Quantum Inform. Comput. **2**, 240 (2002).
 - [17] M.A.Nielsen, Phys. Rev. Lett. , **83**, 436 (1999).